

FINITE TIME BLOW-UP OF SOLUTIONS FOR SYSTEM OF κ TH ORDER WAVE EQUATIONS WITH NONLINEAR AVERAGED DAMPING AND SOURCES

Kh. ZENNIR, K. BOUHALI, H. ZHANG AND D. GUIDAD

Abstract. We consider basic relationships between the initial energy and the exponent of nonlinear sources in a nonlinear coupled system of κ th order wave equations with nonlinear averaged damping, and we demonstrate the global nonexistence of solutions. This approach is a variant of the method of nonlinear functional analysis with a contradiction argument, which is one of the tools for proving the blow-up of solutions for nonlinear partial differential equations. A new class of nonlinear coupled wave equations with nonlinear sources is given in high-order functional spaces.

Keywords: κ th order wave equations, nonlinear equations, coupled system, nonlinear averaged damping, local existence, blow-up.

Mathematics Subject Classification: 35A01; 35L20; 35B44

1. INTRODUCTION, RELEVANCE AND OBJECTIVES

Let $\Omega \subset \mathbb{R}^n$ ($n \geq 1$) be a bounded domain with a smooth boundary Γ . We consider the initial boundary value problem for nonlinear κ th order wave equations with a nonlocal damping

$$\begin{cases} u_{tt} + (-\Delta)^\kappa u + \|u_t\|^l u_t = |u|^{p-2} u |v|^p, \\ v_{tt} + (-\Delta)^\kappa v + \|v_t\|^l v_t = |v|^{p-2} v |u|^p, \end{cases} \quad x \in \Omega, \quad t > 0, \quad (1.1)$$

with boundary conditions

$$\begin{cases} u = \frac{\partial u}{\partial \nu} = \dots = \frac{\partial^{\kappa-1} u}{\partial \nu^{\kappa-1}} = 0, \\ v = \frac{\partial v}{\partial \nu} = \dots = \frac{\partial^{\kappa-1} v}{\partial \nu^{\kappa-1}} = 0, \end{cases} \quad x \in \Gamma, \quad (1.2)$$

and initial conditions

$$\begin{cases} u(x, 0) = u_0(x), & u_t(x, 0) = u_1(x), \\ v(x, 0) = v_0(x), & v_t(x, 0) = v_1(x), \end{cases} \quad x \in \Omega. \quad (1.3)$$

Here ν is the outward normal to the boundary Γ , while $l > 0$, $\kappa \geq 1$ are integers and $p > 2$, and for $u = u(x, t)$

$$\begin{aligned} |\nabla^\kappa u|^2 &= (\Delta^{\frac{\kappa}{2}} u)^2 && \text{for even } \kappa, \\ |\nabla^\kappa u|^2 &= |\nabla(\Delta^{\frac{\kappa-1}{2}} u)|^2 && \text{for even } \kappa, \end{aligned}$$

KH. ZENNIR, K. BOUHALI, H. ZHANG AND D. GUIDAD, FINITE TIME BLOW-UP OF SOLUTIONS FOR SYSTEM OF κ TH ORDER WAVE EQUATIONS WITH NONLINEAR AVERAGED DAMPING AND SOURCES.

© Zennir Kh., Bouhali K., Zhang H. and Guidad D. 2026.

Submitted February 27, 2025.

and

$$|\nabla u|^2 = \sum_{i=1}^n \left(\frac{\partial u}{\partial x_i} \right)^2, \quad \Delta u = \sum_{i=1}^n \frac{\partial^2 u}{\partial x_i^2}.$$

Recently, an initial boundary value problem for following wave equation with nonlinear averaged damping

$$u_{tt} - \Delta u + g\left(\int_{\Omega} |u_t|^2 dx\right)u_t = f(u) \tag{1.4}$$

attracted a lot of attention. Equation (1.4) with the source $f(w) = w \int_{\Omega} |w|^2 dx$ arises from a nonlinear theory of meson fields [6], [11], [14]. It can also be intrinsically connected with the study of damping phenomena in flight structures, where the $\|u_t\|^m$ reflects the effect of kinetic energy; see [2], [7], [19]. For instance, in [3], the one dimensional beam equation

$$u_{tt} - 2\zeta\sqrt{\lambda}u_{xx} + \lambda u_{xxxx} - \gamma \left(\int_L^{-L} (\lambda |u_{xx}|^2 + |u_t|^2 dx) \right)^q u_{xxt} = 0 \tag{1.5}$$

was proposed. As the first stage of the corresponding equation

$$u_{tt} + \left(\int_{\Omega} u_t^2 dx \right)^{\frac{\alpha}{2}} u_t - \Delta u = h(t, x),$$

the pioneering works [1] provided the above equation in a bounded domain Ω with homogeneous boundary conditions; see [1], [5], [17]. Later, the authors in [11] studied the system

$$\begin{aligned} u_{tt} - \mu(t)\Delta u + \alpha f\left(\int_{\Omega} |u|^2 dx\right)u + \beta g\left(\int_{\Omega} |u_t|^2 dx\right)u_t &= 0, \\ u = 0 \quad \text{on} \quad \Gamma_0, \quad \frac{\partial u}{\partial \nu} + h(\cdot, u_t) &= 0 \quad \text{on} \quad \Gamma_1, \\ u(x, 0) = u_0(x), \quad u_t(x, 0) &= u_1(x), \end{aligned} \tag{1.6}$$

under certain conditions on $\mu(t)$, $f(s)$, $g(s)$, $h(s)$. Global existence in time was shown by using the well-known Galerkin method. In addition, a decay rate of the solution was obtained by two different methods, both Lyapunov functional and Nakao’s lemma. Zhang [9] extended and improved the result in [11]. In [22], [23], the long time behavior of the equations

$$u_{tt} + \Delta^2 u - M(\|\nabla u\|^2)\Delta u + k(\|u_t\|)u_t + f(u) = h(x) \tag{1.7}$$

$$u_{tt} - \Delta u + k(\|u_t\|)u_t + f(u) = \int_{\Omega} K(x, y)u_t(y)dy + h(x) \tag{1.8}$$

was considered. The global well-posedness of (1.7) (1.8) was established with the damping $\|u_t\|^m u_t$, $m \geq 0$, see [10], [13], [20] and references therein. In [7], the authors investigated the nonhomogeneous n -dimensional version of (1.5)

$$u_{tt} - \kappa\Delta u + \Delta^2 u - \gamma \left(\int_{\Omega} (|\Delta u|^2 + |u_t|^2) dx \right)^q \Delta u_t + f(u) = 0 \quad \text{in} \quad \Omega \times R^+ \tag{1.9}$$

subject to a boundary condition and with $q \geq 1$. The authors proved the existence of a unique global solution and obtained the polynomial stability and a non-exponential decay prospect of the solutions provided that $q \geq 1$, see [4], [15], [18].

For the equation

$$u_{tt} - \Delta u + \left(\int_{\Omega} |u_t|^2 dx \right)^{\frac{m}{2}} u_t = \left(\int_{\Omega} |u|^p dx \right)^r |u|^{p-2} u \quad (1.10)$$

with homogeneous boundary conditions, Hu, Li, Liu and Zhang [8] provided sufficient conditions for finite time blow-up of weak solutions, with suitable conditions on initial data, by an ordinary differential inequality for an appropriately chosen functional. When the terms $\left(\int_{\Omega} |u|^p dx \right)^r |u|^{p-2} u$ and $-\Delta u$ in equation (1.10) are replaced by $|u|^{p-2} u$ and $\Delta^2 u$, respectively, Zennir and Miyasita [21] proved the global non-existence in time for a large class of pseudo-parabolic equations with weak viscoelasticity under suitable conditions on the variable exponents with negative initial energy.

To our knowledge, there are few results on the non-linear coupled system of κ th order wave equations with nonlinear averaged damping and sources. The main goal of this paper is to study the blow-up conditions of a solution to problem (1.1)–(1.3) with the averaged damping term and the source term, which differ from the existing literature mentioned above. To the best of our knowledge, this is the first result on the nonlinear coupled system of κ th order wave equations with nonlinear averaged damping and sources.

The paper is organized as follows. Section 2 is concerned with some notation and necessary materials. In Section 3, we demonstrate the global nonexistence of solutions with positive or negative initial energy.

2. PRELIMINARIES

Let $L^2(\Omega)$, $H^p(\Omega)$ and $H_0^2(\Omega)$ be the usual Sobolev spaces, where

$$(u, v) = \int_{\Omega} uv dx, \quad \|u\| = \left(\int_{\Omega} |u|^2 dx \right)^{\frac{1}{2}}, \quad \|u\|_p = \left(\int_{\Omega} |u|^p dx \right)^{\frac{1}{p}}.$$

Definition 2.1. *Functions*

$$(u, v) \in C\left([0, T]; H^{\kappa}(\Omega)\right) \cap C^1\left([0, T]; L^2(\Omega)\right) \times C\left([0, T]; H^{\kappa}(\Omega)\right) \cap C^1\left([0, T]; L^2(\Omega)\right),$$

with the initial data given in (1.3) are

1. a strong solution to problem (1.1)–(1.3) on the interval $[0, T]$ if

$$\begin{aligned} (u, v) &\in W^{1,1}(\alpha, \beta; H_0^{\kappa}(\Omega)) \times W^{1,1}(\alpha, \beta; H_0^{\kappa}(\Omega)), \\ (u_t, v_t) &\in W^{1,1}(\alpha, \beta; L^2(\Omega)) \times W^{1,1}(\alpha, \beta; L^2(\Omega)), \end{aligned}$$

for all $0 < \alpha < \beta < T$ and

$$\Delta^{\kappa} u + \Delta^{\kappa} v + \|u_t(t)\|^l u_t + \|v_t(t)\|^l v_t \in L^2(\Omega) \quad \text{for a.e. } t \in [0, T],$$

and (1.1) is satisfied in $L^2(\Omega)$ for all $t \in [0, T]$;

2. a generalized solution to (1.1)–(1.3) on the interval $[0, T]$ if there is a sequence of strong solutions $(\{u_j(t)\}, \{v_j(t)\})$ to (1.1)–(1.3) with initial data $(u_0^j, v_0^j), (u_1^j, v_1^j)$ instead of $(u_0, v_0), (u_1, v_1)$ such that

$$\begin{aligned} (u^j, v^j) &\rightarrow (u, v) \quad \text{in } C([0, T]; H_0^{\kappa}(\Omega)) \times C([0, T]; H_0^{\kappa}(\Omega)) \quad \text{as } j \rightarrow +\infty, \\ (u_t^j, v_t^j) &\rightarrow (u_t, v_t) \quad \text{in } C([0, T]; L^2(\Omega)) \times C([0, T]; L^2(\Omega)) \quad \text{as } j \rightarrow +\infty. \end{aligned}$$

3. a distributed solution to (1.1)–(1.3) on $[0, T]$ if

$$\begin{aligned}
& \int_{\Omega} (u_t(x, t)\phi(x) + v_t(x, t)\psi(x)) dx \\
& + \int_0^t \left(\int_{\Omega} \nabla^{\kappa} u(x, s) \nabla^{\kappa} \phi(x) dx + \|u_t(s)\|^l \int_{\Omega} u_t(x, s)\phi(x) dx \right) ds \\
& + \int_0^t \left(\int_{\Omega} \nabla^{\kappa} v(x, s) \nabla^{\kappa} \psi(x) dx + \|v_t(s)\|^l \int_{\Omega} v_t(x, s)\psi(x) dx \right) ds \\
& = \int_{\Omega} u_1\phi(x) dx + \int_0^t \int_{\Omega} |u|^{p-2}u(x, s)|v(x, s)|^p\phi(x) dx ds \\
& + \int_{\Omega} v_1\psi(x) dx + \int_0^t \int_{\Omega} |v|^{p-2}v(x, s)|u(x, s)|^p\psi(x) dx ds
\end{aligned} \tag{2.1}$$

holds for every test function $\phi, \psi \in H_0^{\kappa}(\Omega) \cap L^2(\Omega)$ and for all $t \in [0, T]$.

Lemma 2.1. [22] For $u_t, w_t \in L^2(\Omega)$, we have

$$\|u_t - w_t\|^{l+2} \leq (\|u_t\|^l u_t - \|w_t\|^l w_t, u_t - w_t). \tag{2.2}$$

The associated energy of (1.1)–(1.3) is defined by

$$E(t) = \frac{1}{2}\|u_t\|^2 + \frac{1}{2}\|v_t\|^2 + J(u, v), \quad 0 \leq t, \tag{2.3}$$

$$J(u, v) = \frac{1}{2}\|\nabla^{\kappa} u\|^2 + \frac{1}{2}\|\nabla^{\kappa} v\|^2 - \frac{1}{p}\|uv\|_p^p, \tag{2.4}$$

and

$$E|_{t=0} = \frac{1}{2}\|u_1\|^2 + \frac{1}{2}\|v_1\|^2 + J(u_0, v_0), \tag{2.5}$$

$$\|u\|_p \leq C_0 \|\nabla^{\kappa} u\|, \quad C_0 > 0, \quad \forall u \in H_0^{\kappa}(\Omega),$$

where

$$\begin{cases} 2 \leq p < +\infty & \text{if } n = \kappa, 2\kappa \\ 2 \leq p \leq \frac{2n}{n-2\kappa} & \text{if } n \geq 3\kappa. \end{cases} \tag{2.6}$$

3. FINITE TIME BLOW UP OF SOLUTIONS

We state the local existence in time of (1.1)–(1.3) by following the lines of [4].

Theorem 3.1. If $(u_0, v_0) \in (H^{2\kappa}(\Omega) \cap H_0^{\kappa}(\Omega)) \times (H^{2\kappa}(\Omega) \cap H_0^{\kappa}(\Omega))$ and $(u_1, v_1) \in H_0^{\kappa}(\Omega) \times H_0^{\kappa}(\Omega)$ are such that

$$\nabla^{\kappa} u_0 + \nabla^{\kappa} v_0 + k\|u_1\|^l u_1 + k\|v_1\|^l v_1 \in L^2(\Omega).$$

Suppose that $l > 0$ and p satisfies

$$\begin{cases} 1 \leq p < +\infty & \text{if } n = \kappa, 2\kappa \\ 1 \leq p \leq \frac{4\kappa - n}{n - 2\kappa} & \text{if } n \geq 3\kappa. \end{cases} \tag{3.1}$$

Then there exists $t_{\max} \leq +\infty$ such that the problem (1.1)–(1.3) has a unique strong solution (u, v) on $[0, t_{\max})$.

If $(u_0, v_0) \in H_0^\kappa(\Omega) \times H_0^\kappa(\Omega)$ and $(u_1, v_1) \in L^2(\Omega) \times L^2(\Omega)$, then there exists $t_{\max} \leq +\infty$ such that (1.1)–(1.3) admits a unique generalized solution, which is also the distributional solution.

With positive and negative initial data, both the strong solution and the weak solution (u, v) cannot be global solutions in $[0, t_{\max})$, $t_{\max} = +\infty$, so for $t_{\max} < \infty$ we have

$$\lim_{t \rightarrow t_{\max}} (\|\nabla^\kappa u\| + \|\nabla^\kappa v\| + \|u_t\| + \|v_t\|) = +\infty, \quad (3.2)$$

The energy relation is satisfied

$$E(t) + \int_0^t (\|u_t(s)\|^{l+2} + \|v_t(s)\|^{l+2}) ds = E(0), \quad 0 \leq t < t_{\max}. \quad (3.3)$$

3.1. Negative initial energy. Our first main result is as follows.

Theorem 3.2. *Let $E(0) < 0$. Under the assumptions of Theorem 3.1 there exists a finite time $t_{\max} < +\infty$, at which the local solution to (1.1)–(1.3) blows up.*

Доказательство. Let

$$\mathcal{K}(t) = -E(t).$$

The contradiction argument will be used. By (2.4) and (3.3) we obtain

$$\begin{aligned} \mathcal{K}'(t) &= -E'(t) \geq 0, \\ 0 < -E(0) = \mathcal{K}(t=0) &\leq \mathcal{K}(t) \leq \frac{1}{p} \|uv\|_p^p. \end{aligned} \quad (3.4)$$

We denote

$$G(t) = \int_{\Omega} (\|u_t\|^l u_t u + \|v_t\|^l v_t v) dx,$$

then, by Hölder and ϖ -Young's inequalities for each $\varpi > 0$ we obtain

$$\begin{aligned} \|u_t\|^2 \|u\|^2 &\leq \varepsilon \|u_t\|^2 + \frac{1}{4\varepsilon} \|u\|^2, \\ |G(t)| &\leq \|u_t\|^l \|u_t\| \|u\| + \|v_t\|^l \|v_t\| \|v\| \\ &\leq \|u_t\|^l \left(\varpi \|u_t\|^2 + \frac{1}{4\varpi} \|u\|^2 \right) + \|v_t\|^l \left(\varpi \|v_t\|^2 + \frac{1}{4\varpi} \|v\|^2 \right) \\ &\leq \varpi [\|u_t\|^{l+2} + \|v_t\|^{l+2}] + \frac{1}{4\varpi} [\|u_t\|^l \|u\|^2 + \|v_t\|^l \|v\|^2] \end{aligned}$$

Again, using Young's inequality with

$$p = \frac{l+2}{l}, \quad q = \frac{l+2}{2}, \quad \frac{1}{p} + \frac{1}{q} = 1,$$

for each $\varpi > 0$ we obtain,

$$\|u_t\|^l \|u\|^2 \leq \varpi \|u_t\|^{lp} + \varpi^{-\frac{2}{l}} \|u\|^{2q} = \varpi \|u_t\|^{l+2} + \varpi^{-\frac{2}{l}} \|u\|^{l+2},$$

and similarly,

$$\|v_t\|^l \|v\|^2 \leq \varpi \|v_t\|^{l+2} + \varpi^{-\frac{2}{l}} \|v\|^{l+2}.$$

Hence,

$$\|u_t\|^l \|u\|^2 + \|v_t\|^l \|v\|^2 \leq \varpi (\|u_t\|^{l+2} + \|v_t\|^{l+2}) + \varpi^{-\frac{2}{l}} (\|u\|^{l+2} + \|v\|^{l+2}),$$

which leads by (3.3) to

$$\begin{aligned} |G(t)| &\leq (\varpi + \frac{1}{4})[\|u_t\|^{l+2} + \|v_t\|^{l+2}] + \frac{\varpi^{-\frac{2+l}{l}}}{4}(\|u\|^{l+2} + \|v\|^{l+2}) \\ &\leq (\varpi + \frac{1}{4})\mathcal{K}'(t) + \frac{\varpi^{-\frac{2+l}{l}}}{4}(\|u\|^{l+2} + \|v\|^{l+2}). \end{aligned} \quad (3.5)$$

We define the functional

$$z(t) = \mathcal{K}^{1-\nu}(t) + \epsilon \int_{\Omega} u_t u \, dx + v_t v \, dx, \quad \epsilon > 0, \quad 0 < \nu < 1. \quad (3.6)$$

Taking the derivative of (3.6) in t , by (1.1), and (3.5), we have

$$\begin{aligned} z'(t) &= (1-\nu)\mathcal{K}^{-\nu}(t)\mathcal{K}'(t) + \epsilon \int_{\Omega} (u_{tt}u + v_{tt}v) \, dx + \epsilon(\|u_t\|^2 + \|v_t\|^2) \\ &= (1-\nu)\mathcal{K}^{-\nu}(t)\mathcal{K}'(t) + \epsilon \left(\|u_t\|^2 + \|v_t\|^2 - \|\nabla^{\kappa}u\|^2 - \|\nabla^{\kappa}v\|^2 + \|uv\|_p^p - G(t) \right) \\ &\geq \left((1-\nu)\mathcal{K}^{-\nu}(t) - \frac{1}{4}\epsilon\varpi \right) \mathcal{K}'(t) + \epsilon \left(\|u_t\|^2 + \|v_t\|^2 - \|\nabla^{\kappa}u\|^2 - \|\nabla^{\kappa}v\|^2 + \|uv\|_p^p \right) \\ &\quad - \frac{\varpi^{-\frac{2+l}{l}}}{4}(\|u\|^{l+2} + \|v\|^{l+2}). \end{aligned} \quad (3.7)$$

We can choose the Young parameter ϖ as a time-dependent weight of $\mathcal{K}(t)$, namely

$$\varpi = k \mathcal{K}^{-\nu}(t), \quad k > 0.$$

Moreover,

$$\varpi^{-\frac{l+2}{l}} = k^{-\frac{l+2}{l}} \mathcal{K}^{\frac{\nu(l+2)}{l}}(t),$$

where the constant k is redefined if necessary.

Substituting these estimates into (3.7), we obtain

$$\begin{aligned} z'(t) &\geq \left(1 - \nu - \epsilon \frac{k}{4} \right) \mathcal{K}^{-\nu}(t) \mathcal{K}'(t) \\ &\quad + \epsilon \left(\|u_t\|^2 + \|v_t\|^2 - \|\nabla^{\kappa}u\|^2 - \|\nabla^{\kappa}v\|^2 + \|uv\|_p^p - \frac{1}{4k^{\frac{l+2}{l}}} \mathcal{K}^{\frac{\nu(l+2)}{l}}(t) (\|u\|^{l+2} + \|v\|^{l+2}) \right), \end{aligned}$$

We take

$$0 \leq \alpha < \frac{p-2}{p}. \quad (3.8)$$

Then,

$$\begin{aligned} z'(t) &\geq \left(1 - \nu - \epsilon \frac{k}{4} \right) \mathcal{K}^{-\nu}(t) \mathcal{K}'(t) + \epsilon p(1-\alpha)\mathcal{K}(t) + \epsilon \left(\frac{p(1-\alpha)}{2} + 1 \right) (\|u_t\|^2 + \|v_t\|^2) \\ &\quad + \epsilon \left(\frac{p(1-\alpha)}{2} - 1 \right) (\|\nabla^{\kappa}u\|^2 + \|\nabla^{\kappa}v\|^2) + \epsilon p \|uv\|_p^p \\ &\quad - \epsilon \frac{1}{4k^{\frac{l+2}{l}}} \mathcal{K}^{\frac{\nu(l+2)}{l}}(t) (\|u\|^{l+2} + \|v\|^{l+2}). \end{aligned} \quad (3.9)$$

As $l > 0$, choosing ν so that

$$0 < \nu < \frac{l}{l+2},$$

and using the upper bound in (3.4), we obtain

$$\mathcal{K}^{\frac{\nu(l+2)}{l}}(t) \leq \mathcal{K}(t) \leq \frac{1}{p} \|uv\|_p^p,$$

and hence

$$\epsilon \frac{1}{4k^{\frac{l+2}{l}}} \mathcal{K}^{\frac{\nu(l+2)}{l}}(t) (\|u\|^{l+2} + \|v\|^{l+2}) \leq \epsilon \frac{1}{4k^{\frac{l+2}{l}} p} \|uv\|_p^p (\|u\|^{l+2} + \|v\|^{l+2}).$$

Since

$$\|u\|^{l+2} + \|v\|^{l+2} \geq 0,$$

it follows immediately that

$$\epsilon \frac{1}{4k^{\frac{l+2}{l}}} \mathcal{K}^{\frac{\nu(l+2)}{l}}(t) (\|u\|^{l+2} + \|v\|^{l+2}) \leq \epsilon \frac{1}{4k^{\frac{l+2}{l}} p} \|uv\|_p^p.$$

Then

$$\begin{aligned} z'(t) \geq & \left(1 - \nu - \epsilon \frac{k}{4}\right) \mathcal{K}^{-\nu}(t) \mathcal{K}'(t) + \epsilon p(1 - \alpha) \mathcal{K}(t) + \epsilon \left(\frac{p(1 - \alpha)}{2} + 1\right) (\|u_t\|^2 + \|v_t\|^2) \\ & + \epsilon \left(\frac{p(1 - \alpha)}{2} - 1\right) (\|\nabla^\kappa u\|^2 + \|\nabla^\kappa v\|^2) + \epsilon \left(p - \frac{1}{4k^{\frac{l+2}{l}} p}\right) \|uv\|_p^p. \end{aligned} \quad (3.10)$$

Choosing α and k such that

$$\begin{aligned} p(1 - \alpha) &> 2, \\ k &> \left(\frac{\epsilon}{4p^2}\right)^{\frac{l}{l+2}}, \end{aligned} \quad (3.11)$$

and then ϵ such that

$$1 - \nu - \epsilon \frac{k}{4} \geq 0,$$

and we have

$$z(t) = \mathcal{K}^{1-\nu}(0) + \epsilon \int_{\Omega} [u_1 u_0 + v_1 v_0] dx > 0.$$

We can pick

$$\nu \in \left(0, \frac{l}{l+2}\right), \quad \alpha \in \left[0, \frac{p-2}{p}\right), \quad k \in \left(\left(\frac{\epsilon}{4p^2}\right)^{\frac{l}{l+2}}, \frac{4(1-\nu)}{\epsilon}\right]$$

so that we can choose parameters satisfying all conditions simultaneously. Then (3.10) becomes

$$z'(t) \geq \epsilon K (\mathcal{K}(t) + \|u_t\|^2 + \|v_t\|^2 + \|uv\|_p^p) \geq 0, \quad K > 0. \quad (3.12)$$

The functional $z(t)$ is increasing on $(0, t_{\max})$ with

$$z(t) \geq z(0) > 0 \quad \text{for all } t \geq 0.$$

We estimate $z^{\frac{1}{1-\nu}}(t)$ by using the elementary inequality

$$|\alpha + \beta|^s \leq 2^{s-1} (|\alpha|^s + |\beta|^s), \quad s \geq 1,$$

and the Young's inequality

$$\begin{aligned} z^{\frac{1}{1-\nu}}(t) &\leq 2^{\frac{1}{1-\nu}} \left(\mathcal{K}(t) + \epsilon \|u\|^{\frac{1}{1-\nu}} \|u_t\|^{\frac{1}{1-\nu}} + \epsilon \|v\|^{\frac{1}{1-\nu}} \|v_t\|^{\frac{1}{1-\nu}} \right) \\ &\leq K \left(\mathcal{K}(t) + \|u_t\|^2 + \|v_t\|^2 + \|u\|^{\frac{2}{1-2\nu}} + \|v\|^{\frac{2}{1-2\nu}} \right). \end{aligned} \quad (3.13)$$

Since $\nu < \frac{1}{2} - \frac{1}{p}$, using the inequality

$$\xi^\tau \leq \left(1 + \frac{1}{s}\right)(s + \xi), \quad \xi \geq 0, \quad 0 \leq \tau \leq 1, \quad s > 0,$$

and taking

$$\xi = \|u\|_2^p, \quad \tau = \frac{2}{(1 - 2\nu)p} < 1,$$

and $s = \mathcal{K}(0)$. We get, for some constants $K_1, K_2 > 0$,

$$\begin{aligned} \|u\|^{\frac{2}{1-2\nu}} + \|v\|^{\frac{2}{1-2\nu}} &\leq \left(1 + \frac{1}{\mathcal{K}(t=0)}\right) (\mathcal{K}(0) + \|u\|_2^p) + \left(1 + \frac{1}{\mathcal{K}(t=0)}\right) (\mathcal{K}(0) + \|v\|_2^p) \\ &\leq K_1 (\mathcal{K}(t) + \|u\|_p^p + \|v\|_p^p) \leq K_2 (\mathcal{K}(t) + \sigma \|uv\|_p^p + C(\sigma)), \quad \sigma > 0. \end{aligned} \tag{3.14}$$

Substituting (3.14) into (3.13), we find

$$z^{\frac{1}{1-\nu}}(t) \leq K_3 (\mathcal{K}(t) + \|u_t\|^2 + \|v_t\|^2 + \|uv\|_p^p), \quad K_3 > 0. \tag{3.15}$$

By (3.12), (3.15), we get

$$z'(t) \geq K_4 z^{\frac{1}{1-\nu}}(t), \quad K_4 > 0. \tag{3.16}$$

A simple integration of (3.16) over $(0, t)$ then yields

$$z(t) \geq \left(z^{\frac{-\nu}{1-\nu}}(t=0) - K_4 \frac{\nu t}{1-\nu} \right)^{\frac{\nu-1}{\nu}}.$$

Thus, there exists t_2 such that

$$\begin{aligned} t_2 \leq t_* &= \frac{z^{\frac{-\nu}{1-\nu}}(t=0)(1-\nu)}{K_4 \nu}, \\ \lim_{t \rightarrow t_2^-} z(t) &= +\infty. \end{aligned} \tag{3.17}$$

The proof is complete. □

3.2. Positive initial energy. Let

$$\begin{aligned} 0 &\leq E(t=0) \leq E_1, \\ E_1 &= \left(\frac{1}{2} - \frac{1}{p}\right) C_0^{-\frac{2p}{p-2}}, \end{aligned} \tag{3.18}$$

and C_0 is the best Poincaré constant.

Lemma 3.1. [16] *If*

$$0 \leq E(t=0) < E_1, \quad \|\nabla^\kappa u_0\| + \|\nabla^\kappa v_0\| > \lambda_1,$$

then there exists $\lambda_2 > \lambda_1$ such that

$$\|\nabla^\kappa u\| + \|\nabla^\kappa v\| \geq \lambda_2, \quad \lambda_1 = C_0^{-\frac{p}{p-2}}.$$

Theorem 3.3. *Under the assumptions of Theorem 3.1, for*

$$0 \leq E(0) < E_1, \quad \|\nabla^\kappa u_0\| + \|\nabla^\kappa v_0\| > \lambda_1,$$

the solution to problem (1.1)-(1.3) blows up, that is, $t_{\max} < +\infty$.

Доказательство. We set

$$\mathcal{K}(t) = E_2 - E(t), \quad E_2 = \frac{E(t=0) + E_1}{2}.$$

Proceeding as in (3.9) with $\alpha = 0$, we obtain

$$\begin{aligned} z'(t) &\geq \left(1 - \nu - \epsilon \frac{k}{4}\right) \mathcal{K}^{-\nu}(t) \mathcal{K}'(t) + \epsilon p \mathcal{K}(t) + \epsilon \left(\frac{p}{2} + 1\right) (\|u_t\|^2 + \|v_t\|^2) \\ &\quad + \epsilon \left(\frac{p}{2} - 1\right) (\|\nabla^\kappa u\|^2 + \|\nabla^\kappa v\|^2) + \epsilon p \|uv\|_p^p \\ &\quad - \frac{\epsilon}{4k^{\frac{l+2}{l}}} \mathcal{K}^{\frac{\nu(l+2)}{l}}(t) (\|u\|^{l+2} + \|v\|^{l+2}) - \epsilon p E_2. \end{aligned} \tag{3.19}$$

Moreover,

$$\begin{aligned} \epsilon \left(\frac{p}{2} - 1\right) (\|\nabla^\kappa u\|^2 + \|\nabla^\kappa v\|^2) - \epsilon p E_2 &= \epsilon \left(\left(\frac{p}{2} - 1\right) (\|\nabla^\kappa u\|^2 + \|\nabla^\kappa v\|^2) - p E_2 \right) \\ &= \epsilon \left(\left(\frac{p}{2} - 1\right) \frac{\lambda_2^2 - \lambda_1^2}{\lambda_2^2} (\|\nabla^\kappa u\|^2 + \|\nabla^\kappa v\|^2) \right. \\ &\quad \left. + \left(\frac{p}{2} - 1\right) \lambda_1^2 \frac{\|\nabla^\kappa u\|^2 + \|\nabla^\kappa v\|^2}{\lambda_2^2} - p E_2 \right) \\ &\geq \epsilon (K_5 (\|\nabla^\kappa u\|^2 + \|\nabla^\kappa v\|^2) + K_6), \end{aligned} \tag{3.20}$$

where λ_2 is given in Lemma 3.1,

$$K_5 = \left(\frac{p}{2} - 1\right) \frac{\lambda_2^2 - \lambda_1^2}{\lambda_2^2} > 0, \quad K_6 = \left(\frac{p}{2} - 1\right) \lambda_1^2 - p E_2.$$

By (3.18) and Lemma 3.1, we observe that

$$K_6 = \left(\frac{p}{2} - 1\right) \lambda_1^2 - \frac{p(E_1 + E(t=0))}{2} = \frac{p(E_1 - E(t=0))}{2} > 0. \tag{3.21}$$

Thus, by (3.20), (3.21) and (3.19), we arrive at

$$\begin{aligned} z'(t) &\geq \left(1 - \nu - \epsilon \frac{k}{4}\right) \mathcal{K}^{-\nu}(t) \mathcal{K}'(t) + \epsilon p \mathcal{K}(t) + \epsilon \left(\frac{p}{2} + 1\right) (\|u_t\|^2 + \|v_t\|^2) \\ &\quad + \epsilon K_5 (\|\nabla^\kappa u\|^2 + \|\nabla^\kappa v\|^2) + \epsilon p \|uv\|_p^p - \epsilon \frac{1}{4k^{\frac{l+2}{l}}} \mathcal{K}^{\frac{\nu(l+2)}{l}}(t) (\|u\|^{l+2} + \|v\|^{l+2}). \end{aligned}$$

Similarly to (3.14) we have

$$\begin{aligned} z'(t) &\geq \left(1 - \nu - \epsilon \frac{k}{4}\right) \mathcal{K}^{-\nu}(t) \mathcal{K}'(t) + \epsilon p \mathcal{K}(t) + \epsilon \left(\frac{p}{2} + 1\right) (\|u_t\|^2 + \|v_t\|^2) \\ &\quad + \epsilon K_5 (\|\nabla^\kappa u\|^2 + \|\nabla^\kappa v\|^2) + \epsilon \left(p - \epsilon \frac{\sigma}{4k^{\frac{l+2}{l}} p}\right) \|uv\|_p^p + C(\sigma). \end{aligned}$$

Choosing k as in (3.11), we can complete the proof as in Theorem 3.2. □

REFERENCES

1. F. Aloui, I.B. Hassen, A. Haraux. *Compactness of trajectories to some nonlinear second order evolution equations and applications* // J. Math. Pures Appl. **100**:3, 295–326 (2013). <https://doi.org/10.1016/j.matpur.2013.01.002>

2. S. F. Aljurbua, A. Bchatnia, A. Beniani, A. Safsaf. *Blow-up for coupled nonlinear wave equations with fractional damping and logarithmic source terms* // Comput. Appl. Math. **44**:8, 400 (2025). <https://doi.org/10.1007/s40314-025-03362-0>
3. A.V. Balakrishnan and L.W. Taylor. *Distributed parameter nonlinear damping models for flight structures* // in Proceedings “Damping 89”, Flight Dynamics Lab and Air Force Wright Aeronautical Labs, WPAFB (1989).
4. I. Chueshov and I. Lasiecka. *Long-Time Behavior of Second Order Evolution Equations with Nonlinear Damping*. Amer. Math. Soc., Providence, Rhode Island (2008). <https://doi.org/10.1090/memo/0912>
5. M. Ghisi, M. Gobbino and A. Haraux. *Quantization of energy and weakly turbulent profiles of solutions to some damped second-order evolution equations* // Adv. Nonlinear Anal. **8**, 902–927 (2019). <https://doi.org/10.1515/anona-2017-0181>
6. K. Jörgens. *Des aufangswert in grossen für eine klasse nichtlinearer wallengleichungen* // Math. Z. **77**:1, 295–308 (1961). <https://doi.org/10.1007/BF01180181>
7. M.A. Jorge Silva, V. Narciso, and A. Vicente. *On a beam model related to flight structures with nonlocal energy damping* // Discrete Contin. Dyn. Syst., Ser. B **24**:7, 3281–3298 (2019). <https://doi.org/10.3934/dcdsb.2018320>
8. Q. Y. Hu, D. H. Li, S. Liu, H. W. Zhang. *Blow-up of solutions for a wave equation with nonlinear averaged damping and nonlocal nonlinear source terms* // Quaest. Math. **46**:4, 695–710 (2023). <https://doi.org/10.2989/16073606.2022.2039972>
9. D. H. Li, H. W. Zhang, Q. Y. Hu. *General energy decay of solutions for a wave equation with nonlocal damping and nonlinear boundary damping* // J. Part. Diff. Equ. **32**:4, 369–380 (2019). <https://doi.org/10.4208/jpde.v32.n4.6>
10. D.H. Li, H.W. Zhang, S. Liu, and Q.Y. Hu. *Blow-up of solutions to a viscoelastic wave equation with nonlocal damping* // Evol. Equ. Control Theory **11**:6, 2017–2031 (2022). <https://doi.org/10.3934/eect.2022009>
11. A. T. Lourêdo, M. A. Ferreira de Araújo, M. M. Miranda. *On a nonlinear wave equation with boundary damping* // Math. Meth. Appl. Sci. **37**:9, 1278–1302 (2014). <https://doi.org/10.1002/mma.2885>
12. S. A. Messaoudi. *Global existence and nonexistence in a system of Petrovsky* // J. Math. Anal. Appl. **265**:2, 296–308 (2002). <https://doi.org/10.1006/jmaa.2001.7697>
13. Q.Q. Peng and Z.F. Zhang. *Global attractor for a coupled wave and plate equation with nonlocal weak damping on Riemannian manifolds* // Appl. Math. Optim. **88**:2, 28 (2023). <https://doi.org/10.1007/s00245-023-09998-w>
14. L. Schiff. *Nonlinear meson theory of nuclear forces I. Neutral scalar mesons with point-contact repulsion* // Phys. Reviews. **84**:1, 1–9 (1951). <https://doi.org/10.1103/PhysRev.84.1>
15. Y. Sun, Z. J. Yang. *Attractors and their continuity for an extensible beam equation with rotational inertia and nonlocal energy damping* // J. Math. Anal. Appl. **512**:2, 126148 (2022). <https://doi.org/10.1016/j.jmaa.2022.126148>
16. S.T. Wu and L.Y. Tsai. *On global solutions and blow-up of solutions for a nonlinearly damped Petrovsky system* // Taiwanese J. Math. **19**:2A, 545–558 (2009). <https://doi.org/10.11650/twjmath/1500405355>
17. Kh. Zennir, M. Bayoud, and S. Georgiev. *Decay of solution for degenerate wave equation of Kirchhoff type in viscoelasticity* // Int. J. Appl. Comput. Math. **4**:1, 54 (2018). <https://doi.org/10.1007/s40819-018-0488-8>
18. Kh. Zennir, A. Guesmia. *Existence of solutions to nonlinear κ th-order coupled Klein — Gordon equations with nonlinear sources and memory terms* // Appl. Math. E-Notes. **15**, 121–136 (2015). <https://emis.de/ft/8211>
19. H. Zhang, L. Liu, H. Yue, D. Li, Kh. Zennir. *The existence, asymptotic behaviour and blow-up of solution of a plate equation with nonlinear averaged damping* // Rep. Math. Phys. **19**:3, 305–323 (2024). [https://doi.org/10.1016/S0034-4877\(24\)00081-8](https://doi.org/10.1016/S0034-4877(24)00081-8)

20. H.W. Zhang, D.H. Li, W.X. Zhang, and Q.Y. Hu. *Asymptotic stability and blow-up for the wave equation with degenerate nonlocal nonlinear damping and source terms* // Appl. Anal. **101**:9, 3170–3181 (2022). <https://doi.org/10.1080/00036811.2020.1836354>
21. Kh. Zennir, T. Miyasita. *Lifespan of solutions for a class of pseudo-parabolic equation with weak-memory* // Alexandria Eng. J. **59**:2, 957–964 (2020). <https://doi.org/10.1016/j.aej.2020.03.016>
22. C.Y. Zhao, C.X. Zhao, C.K.Zhong. *The global attractor for a class of extensible beams with nonlocal weak damping* // Discrete Contin. Dyn. Syst., Ser. B **25**:3, 935–955 (2020). <https://doi.org/10.3934/dcdsb.2019197>
23. C.X. Zhao, F.J. Meng. *Attractor for the extensible beam equation with nonlocal weak damping on time-dependent space* // Acta Math. Sin., Engl. Ser. **40**:4, 1115–1126 (2024). <https://doi.org/10.1007/s10114-023-1295-7>

Khaled Zennir,
Department of Mathematics,
College of Science,
Qassim University,
Buraidah, Saudi Arabia.
E-mail: k.zennir@qu.edu.sa

Keltoum Bouhali,
Department of Mathematics,
College of Science,
Qassim University,
Buraidah, Saudi Arabia.
E-mail: k.bouhali@qu.edu.sa

Hongwei Zhang,
School of Mathematics and Statistics,
Henan University of Technology,
Zhengzhou 450001, China.
E-mail: whz661@163.com

Derradji Guidad,
Department of Mathematics,
College of Sciences,
University of Biskra,
BP 145 RP, 07000 Biskra, Algeria.
E-mail: guidadd07@gmail.com